

# Geometrical Superconformal Anomalies

Johanna Erdmenger<sup>1</sup> and Christian Rupp<sup>2</sup>

Institut für Theoretische Physik  
Universität Leipzig  
Augustusplatz 10/11  
D - 04109 Leipzig  
Germany

## Abstract

We determine the structure of geometrical superconformal anomalies for  $N = 1$  supersymmetric quantum field theories on curved superspace to all orders in  $\hbar$ . For the massless Wess-Zumino model we show how these anomalies contribute to the local Callan-Symanzik equation which expresses the breakdown of superconformal symmetry in terms of the usual  $\beta$  and  $\gamma$  functions.

PACS numbers: 04.62+v, 11.10Gh, 11.10Hi, 11.30Pb.

Keywords: Quantum Field Theory, Superconformal Symmetry,  
Curved Superspace Background, Anomalies.

---

<sup>1</sup>Supported by Deutsche Forschungsgemeinschaft, e-mail: Johanna.Erdmenger@itp.uni-leipzig.de

<sup>2</sup>Supported by Graduiertenkolleg "Quantenfeldtheorie: Mathematische Struktur und physikalische Anwendungen", e-mail: Christian.Rupp@itp.uni-leipzig.de

Superconformal symmetry transformations of a given  $N = 1$  supersymmetric quantum field theory are conveniently studied within the superfield formalism by coupling the theory to a classical minimal supergravity background. For diffeomorphism invariant models the superconformal transformations are then determined by the super Weyl transformations.

We consider curved space backgrounds characterised by a Weyl invariant supergravity prepotential  $H_{\alpha\dot{\alpha}}$ , which couples to the supercurrent of the theory, and a chiral compensator  $\phi \equiv e^J$  whose Weyl transformation is given by  $\delta_\sigma J = \sigma$  with  $\sigma$  chiral.

For the four-dimensional massless Wess-Zumino model on curved superspace, whose dynamical fields are denoted by  $A, \bar{A}$  and the dynamical coupling by  $g$ , it was shown in [1] in a perturbative approach that the *local Callan-Symanzik equation*

$$\left( w^{(\sigma)} - \gamma A \frac{\delta}{\delta A} \right) \Gamma = [\beta^g \partial_g \mathcal{L}_{\text{eff}}] \cdot \Gamma + C(H, J, \bar{J}), \quad (1)$$

$$w^{(\sigma)} = \frac{\delta}{\delta J} - A \frac{\delta}{\delta A}, \quad (2)$$

holds to all orders in  $\hbar$ . Here  $w^{(\sigma)}$  is the chiral Weyl symmetry operator,  $\Gamma$  the vertex functional and  $\beta^g$  and  $\gamma$  are the well-known functions expressing the anomalies of scale invariance. The first term on the right hand side is an insertion which is uniquely determined in terms of field polynomials involving the dynamical fields.  $C(H, J, \bar{J})$  denotes all purely geometrical superconformal anomalies. The aim of this letter is to determine  $C(H, J, \bar{J})$  explicitly to all orders in  $\hbar$ .<sup>3</sup>

A basis for the geometrical terms  $C(H, J, \bar{J})$  in the local Callan-Symanzik equation (1) is given by

$$\begin{aligned} \mathcal{L}_{\text{top}} &= \phi^3 (W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} - (\bar{\mathcal{D}}^2 + R)(G^a G_a - 2R\bar{R})) \\ \mathcal{L}_{\text{Weyl}} &= \phi^3 W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \\ \mathcal{L}_{R\bar{R}} &= \phi^3 (\bar{\mathcal{D}}^2 + R)(R\bar{R}) \\ \mathcal{L}_3 &= \phi^3 R^3 \\ \mathcal{L}'_3 &= \phi^3 (\bar{\mathcal{D}}^2 + R)\bar{R}^2 \\ \mathcal{L}_4 &= \phi^3 (\bar{\mathcal{D}}^2 + R)\mathcal{D}^2 R \equiv \phi^3 \square R. \end{aligned} \quad (3)$$

$W_{\alpha\beta\gamma}$ ,  $G_a$  and  $R$  are the supersymmetric generalisations of the Weyl tensor, Ricci tensor and Ricci scalar respectively.  $G_a$  is a real vector superfield, while  $W_{\alpha\beta\gamma}$  and  $R$  are chiral with their antichiral partners given by  $\bar{W}_{\dot{\alpha}\dot{\beta}\dot{\gamma}}$ ,  $\bar{R}$ . These structures are discussed further in textbooks like [2]. The supercovariant derivatives are denoted by  $\mathcal{D}$ ,  $\bar{\mathcal{D}}$ ;  $(\bar{\mathcal{D}}^2 + R)$  is

---

<sup>3</sup>Our conventions are those of [1].

the chiral projection operator. For the integrals of the basis (3) we define

$$\begin{aligned}
\int d^6 z \mathcal{L}_{\text{top}} &= I_{\text{top}} & \int d^6 z \mathcal{L}_{\text{Weyl}} &= I_{\text{Weyl}} \\
\int d^6 z \mathcal{L}_{R\bar{R}} &= I_{R\bar{R}} & \int d^6 z \mathcal{L}_4 &= I_4 \\
\int d^6 z \mathcal{L}_3 &= I_3 & \int d^6 z \mathcal{L}'_3 &= \bar{I}_3,
\end{aligned} \tag{4}$$

such that we obtain

$$\begin{aligned}
I_{\text{top}} &= \int d^6 z \phi^3 W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} - \int d^8 z E^{-1} (G^a G_a - 2R\bar{R}) \\
I_{\text{Weyl}} &= \int d^6 z \phi^3 W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \\
I_{R\bar{R}} &= \int d^8 z E^{-1} R\bar{R} \\
I_3 &= \int d^6 z \phi^3 R^3 \\
\bar{I}_3 &= \int d^6 \bar{z} \bar{\phi}^3 \bar{R}^3 \\
I_4 &= 0,
\end{aligned} \tag{5}$$

with  $\int d^8 z E^{-1}$  the integration measure for curved superspace.  $\mathcal{L}_{\text{top}}$  and  $\mathcal{L}_{\text{Weyl}}$  are the supersymmetric Gauß-Bonnet and Weyl densities, such that in an expansion around flat space we have

$$I_{\text{Weyl}} - \bar{I}_{\text{Weyl}} = I_{\text{top}} - \bar{I}_{\text{top}} = 0, \tag{6a}$$

$$I_{\text{top}} + \bar{I}_{\text{top}} = 0. \tag{6b}$$

The basis (3) has been investigated using algebraic renormalisation in [3] to first order in  $\hbar$ . In [3] it was found that only the Gauß-Bonnet and Weyl terms are anomalies, while  $\mathcal{L}_3$ ,  $\mathcal{L}'_3$  and  $\mathcal{L}_4$  may be removed by suitable redefinitions of the vertex functional, and  $\mathcal{L}_{R\bar{R}}$  is eliminated by Wess-Zumino consistency. Here we extend this analysis to all orders in  $\hbar$  and discuss the consequences for the local Callan-Symanzik equation (1).

For the Weyl transformation properties of the geometrical terms we have

$$\begin{aligned}
w^{(\sigma)} I_3 &= -3\mathcal{L}_3 & \bar{w}^{(\bar{\sigma})} I_3 &= 3\bar{\mathcal{L}}'_3 \\
w^{(\sigma)} \bar{I}_3 &= 3\mathcal{L}'_3 & \bar{w}^{(\bar{\sigma})} \bar{I}_3 &= -3\bar{\mathcal{L}}_3 \\
w^{(\sigma)} I_{R\bar{R}} &= \mathcal{L}_4 & \bar{w}^{(\bar{\sigma})} I_{R\bar{R}} &= \bar{\mathcal{L}}_4 \\
w^{(\sigma)} I_{\text{Weyl}} &= 0 & \bar{w}^{(\bar{\sigma})} I_{\text{Weyl}} &= 0 \\
w^{(\sigma)} I_{\text{top}} &= 0 & \bar{w}^{(\bar{\sigma})} I_{\text{top}} &= 0.
\end{aligned} \tag{7}$$

By inserting the Weyl transformation parameters  $\sigma$ ,  $\bar{\sigma}$  appropriate for  $R$  transformations and dilatations [1] into the Weyl operator

$$\mathcal{W} = -i \int d^6 z \sigma w^{(\sigma)} - i \int d^6 \bar{z} \bar{\sigma} \bar{w}^{(\bar{\sigma})},$$

we obtain global Ward operators

$$\mathcal{W}^R = \frac{2}{3} \int d^6 z w^{(\sigma)} - \frac{2}{3} \int d^6 \bar{z} \bar{w}^{(\bar{\sigma})}, \quad (8)$$

$$\mathcal{W}^D = i \int d^6 z w^{(\sigma)} + i \int d^6 \bar{z} \bar{w}^{(\bar{\sigma})} \quad (9)$$

for global R transformations and dilatations respectively. By applying these operators to the integrated expressions (5), we find that all these expressions are invariant under R transformations and dilatations except  $I_3$  and  $\bar{I}_3$ , for which

$$\mathcal{W}^R(I_3 + \bar{I}_3) = -4(I_3 - \bar{I}_3), \quad \mathcal{W}^D I_3 = 0, \quad \mathcal{W}^D \bar{I}_3 = 0. \quad (10)$$

Within the BPHZ approach pursued here [4], the renormalisation process necessitates the introduction of an auxiliary mass term  $M(s-1) (\int d^6 z \phi^3 A^2 + \int d^6 \bar{z} \bar{\phi}^3 \bar{A}^2)$  even within the massless theory [5]. This term is at the origin of potential anomalies since it leads to symmetry breaking hard insertions involving

$$[M(s-1)\phi^3 A^2] \cdot \Gamma. \quad (11)$$

Additional information is needed to evaluate such terms before the limit  $s = 1$  may be taken since the parameter  $s$  participates in the subtraction process similarly to an external momentum. This information is provided by the Zimmermann identities [6] according to which terms like (11) may be expressed by a basis of local field polynomial insertions. In our case this leads to

$$[M(s-1)\phi^3 A^2] \cdot \Gamma|_{s=1} = [Z_{\text{dyn}}] \cdot \Gamma + Z_{\text{geom}}, \quad (12)$$

where the terms  $Z_{\text{dyn}}$  involving dynamical fields have been discussed in [1] and the geometrical contributions  $Z_{\text{geom}}$  are given by

$$Z_{\text{geom}} = u_{\text{top}} \mathcal{L}_{\text{top}} + u_{\text{Weyl}} \mathcal{L}_{\text{Weyl}} + u_{R\bar{R}} \mathcal{L}_{R\bar{R}} + u_3 \mathcal{L}_3 + u'_3 \mathcal{L}'_3 + u_4 \mathcal{L}_4 \quad (13)$$

in terms of the basis (3). The coefficients  $u$  are power series in  $\hbar$ .

In the BPHZ approach, vertex functions are calculated from the *effective action*  $\Gamma_{\text{eff}}$  in agreement with the forest formula. Here we have

$$\Gamma_{\text{eff}} = \Gamma_{\text{eff}}^{\text{dyn}} + \Gamma_{\text{eff}}^{\text{geom}}. \quad (14)$$

We specify the geometrical contribution  $\Gamma_{\text{eff}}^{\text{geom}}$  in such a way that the removable symmetry breaking terms arising from the Zimmermann identity may be cancelled,

$$\Gamma_{\text{eff}}^{\text{geom}} = \frac{1}{8} \lambda_3 (I_3 + \bar{I}_3) + \frac{1}{8} \lambda_{R\bar{R}} I_{R\bar{R}}. \quad (15)$$

With these results we obtain for the Weyl transformation of the Wess-Zumino vertex functional  $\Gamma$  on curved superspace using (7) and (12)

$$w^{(\sigma)} \Gamma|_{s=1} = -\frac{3}{2} [S_{\text{dyn}}] \cdot \Gamma|_{s=1} - \frac{3}{2} S_{\text{geom}} \quad (16)$$

with  $S_{\text{dyn}}$  as determined in [1] and<sup>4</sup>

$$S_{\text{geom}} = \frac{1}{12} \left( u_{\text{top}} \mathcal{L}_{\text{top}} + u_{\text{Weyl}} \mathcal{L}_{\text{Weyl}} + u_{R\bar{R}} \mathcal{L}_{R\bar{R}} + (u_3 + 3\lambda_3) \mathcal{L}_3 + (u'_3 - 3\lambda_3) \mathcal{L}'_3 + (u_4 - \lambda_{R\bar{R}}) \mathcal{L}_4 \right). \quad (17)$$

Due to (6a) the only terms in (16) contributing to global  $R$  transformations (8) are

$$\mathcal{W}^R \Gamma \Big|_{s=1} = -\frac{1}{12} (u_3 - u'_3 + 6\lambda_3) (I_3 - \bar{I}_3), \quad (18)$$

such that when setting

$$\lambda_3 = \frac{1}{6} (u'_3 - u_3) \quad (19)$$

we have global  $R$  invariance. At the local level however,  $R$  symmetry is anomalous since the non-integrated terms (17) and their antichiral partners contribute to the divergence of the supercurrent.

The global Callan-Symanzik equation obtained by applying the operator  $\mathcal{C} \equiv m\partial_m + \beta^g \partial_g - \gamma \mathcal{N}$  to  $\Gamma$ , with  $\mathcal{N} = \int d^6 z A \frac{\delta}{\delta A} + \int d^6 \bar{z} \bar{A} \frac{\delta}{\delta \bar{A}}$  the operator counting matter legs, reads

$$\mathcal{C} \Gamma \Big|_{s=1, H=J=0} = 0 \quad (20)$$

on flat space. This equation acquires anomalies on curved superspace as well, which are of the form

$$\mathcal{C} \Gamma \Big|_{s=1} = \Delta_{\mathcal{C}}^{\text{geom}}, \quad (21)$$

$$\begin{aligned} \Delta_{\mathcal{C}}^{\text{geom}} = & -\frac{1}{8} (1 - 2\gamma) u_{\text{Weyl}} (I_{\text{Weyl}} + \bar{I}_{\text{Weyl}}) + \frac{1}{8} (\beta^g \partial_g \lambda_{R\bar{R}} - 2(1 - 2\gamma) u_{R\bar{R}}) I_{R\bar{R}} \\ & + \frac{1}{8} (\beta^g \partial_g \lambda_3 - (1 - 2\gamma)(u_3 + u'_3)) (I_3 + \bar{I}_3). \end{aligned} \quad (22)$$

Note that due to (6b) the Gauß-Bonnet term does not contribute to the anomalies of the global Callan-Symanzik equation in agreement with the topological nature of this term from which follows its scale independence. Furthermore, using the consistency condition  $[\mathcal{W}^R, \mathcal{C}] \Gamma = 0$  we may show that the coefficient of the  $I_3 + \bar{I}_3$  term in (22) vanishes,

$$\beta^g \partial_g \lambda_3 - (1 - 2\gamma)(u_3 + u'_3) = 0, \quad (23)$$

such that (22) reduces to

$$\Delta_{\mathcal{C}}^{\text{geom}} = -\frac{1}{8} (1 - 2\gamma) u_{\text{Weyl}} (I_{\text{Weyl}} + \bar{I}_{\text{Weyl}}) + \frac{1}{8} (\beta^g \partial_g \lambda_{R\bar{R}} - 2(1 - 2\gamma) u_{R\bar{R}}) I_{R\bar{R}}. \quad (24)$$

---

<sup>4</sup>To first order in  $\hbar$  the coefficients  $u$  in (17) have been calculated explicitly in a different approach [7].

Similarly we determine the geometrical terms  $C(H, J, \bar{J})$  in the local Callan-Symanzik equation (1). For the R invariant theory (19), they are given by

$$\left( w^{(\sigma)} - \gamma A \frac{\delta}{\delta A} \right) \Gamma \Big|_{s=1} = -\beta^g [\partial_g \mathcal{L}_{\text{eff}}] \cdot \Gamma \Big|_{s=1} + C(H, J, \bar{J}), \quad (25)$$

$$\begin{aligned} C(H, J, \bar{J}) = & -\frac{1}{8}(1-2\gamma)u_{\text{top}}\mathcal{L}_{\text{top}} - \frac{1}{8}(1-2\gamma)u_{\text{Weyl}}\mathcal{L}_{\text{Weyl}} \\ & + \frac{1}{8} \left( \frac{1}{2}\beta^g \partial_g \lambda_{R\bar{R}} - (1-2\gamma)u_{R\bar{R}} \right) \mathcal{L}_{R\bar{R}} \\ & + \frac{1}{8} \left( \frac{1}{2}(u_3 + u'_3) - 2\gamma u'_3 \right) (\mathcal{L}_3 - \mathcal{L}'_3), \end{aligned} \quad (26)$$

with the geometrical part of  $\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{dyn}} + \mathcal{L}_{\text{geom}}$  given by

$$\mathcal{L}_{\text{geom}} \equiv \frac{1}{8}\lambda_3\mathcal{L}_3 + \frac{1}{16}\lambda_{R\bar{R}}\mathcal{L}_{R\bar{R}}. \quad (27)$$

Here we have chosen  $\lambda_{R\bar{R}} = (1-2\gamma)u_4$  in order to absorb the breaking term involving  $\mathcal{L}_4$ .

Integrating (25) over chiral superspace and adding the complex conjugate we recover the global Callan-Symanzik equation (21) by virtue of (9) and of  $\mathcal{W}^D \Gamma = im \partial_m \Gamma$ . The coefficients of the superconformal anomalies given by  $C(H, J, \bar{J})$  are in general non-zero to any given order in  $\hbar$ . However there are simplifications in special cases. To first order in  $\hbar$  the consistency equation

$$[w^{(\sigma)}(z), \bar{w}^{(\bar{\sigma})}(z')] \Gamma = 0 \quad (28)$$

implies

$$u_3 + u'_3 = O(\hbar^2), \quad u_{R\bar{R}} = O(\hbar^2), \quad (29)$$

such that we obtain

$$\left( w^{(\sigma)} - \gamma A \frac{\delta}{\delta A} \right) \Gamma \Big|_{s=1} = -\beta^g [\partial_g \mathcal{L}_{\text{eff}}] \cdot \Gamma \Big|_{s=1} - \frac{1}{8}u_{\text{top}}\mathcal{L}_{\text{top}} - \frac{1}{8}u_{\text{Weyl}}\mathcal{L}_{\text{Weyl}} + O(\hbar^2). \quad (30)$$

Furthermore at fixed points where  $\beta^g = 0$  it is straightforward to determine the consequences of the consistency condition (28) to all orders in  $\hbar$  since no double insertions appear when applying  $w^{(\sigma)}$  to the r.h.s. of (25) in this case. Therefore for  $\beta^g = 0$  the relations (29) are valid to all orders in  $\hbar$ , and (25) simplifies to

$$\begin{aligned} \left( w^{(\sigma)} - \gamma A \frac{\delta}{\delta A} \right) \Gamma \Big|_{s=1} = & -\frac{1}{8}(1-2\gamma)(u_{\text{top}}\mathcal{L}_{\text{top}} + u_{\text{Weyl}}\mathcal{L}_{\text{Weyl}}) \\ & - \frac{1}{4}\gamma u'_3(\mathcal{L}_3 - \mathcal{L}'_3), \end{aligned} \quad (31)$$

with  $\gamma u'_3 = O(\hbar^2)$ . For the trivial fixed point of the Wess-Zumino model where  $g = \gamma = 0$  this reduces to

$$w^{(\sigma)} \Gamma \Big|_{s=1} = -\frac{1}{8}u_{\text{top}}\mathcal{L}_{\text{top}} - \frac{1}{8}u_{\text{Weyl}}\mathcal{L}_{\text{Weyl}}. \quad (32)$$

For the Wess-Zumino model this confirms that at the fixed point, the structure of correlation functions involving the supercurrent is entirely determined by (32), as was recently discussed in [8].

# References

- [1] J. Erdmenger, C. Rupp and K. Sibold, *Conformal Transformation Properties of the Supercurrent in Four Dimensional Supersymmetric Theories*, Nucl. Phys. **B** to be published, hep-th/9804053.
- [2] I.L. Buchbinder and S.M. Kuzenko, *Ideas and Methods of Supersymmetry and Supergravity*. Institute of Physics Publishing, Bristol and Philadelphia 1995.
- [3] L. Bonora, P. Pasti and M. Tonin, Nucl.Phys. **B 252** (1985) 458.
- [4] J.H. Lowenstein, *BPHZ renormalization*, in: Renormalization Theory, eds. G. Velo and A.S. Wightman, Erice lectures 1976;  
O. Piguet and S. Sorella, *Algebraic Renormalization*. Springer Verlag, Berlin 1995;  
J. Collins, *Renormalization*. Cambridge University Press 1984.
- [5] W. Zimmermann, Commun.Math.Phys **15** (1969) 208;  
J.H. Lowenstein and W. Zimmermann, Commun.Math.Phys. **44** (1975) 73;  
J.H. Lowenstein, Commun.Math.Phys. **47** (1976) 53.
- [6] W. Zimmermann, Ann.Phys. (NY) **77** (1973) 536;  
T.E. Clark and J.H. Lowenstein, Nucl.Phys. **B 113** (1976) 109.
- [7] I.N. McArthur, Class. Quant. Grav. **1** (1984) 245;  
I.L. Buchbinder and S.M. Kuzenko, Nucl. Phys. **B 274** (1986) 653.
- [8] H. Osborn,  *$N = 1$  Superconformal Symmetry in Four Dimensional Quantum Field Theory*, hep-th/9808041.